## Applied PhytoGenetics, Inc.

- ➢APGEN, Inc. is a phytoremediation company.
- ➢Phytoremediation is using plants to clean polluted soil and water
  - Degrade organics
  - Detoxify and sequester toxic elements like heavy metals



APGEN, Inc 12/8/00 1

#### Scientific Team

Richard B. Meagher, Ph.D. - Chief Scientific Officer

**Laura Carreira**, Ph.D. - Principal Research Officer



APGEN, Inc 12/8/00 2

## APGEN is based in Georgia

- **≈**APGEN is located in University of Georgia's Biotechnology Development space
  - Rm 169, 110 Riverbend Rd.
  - Athens, GA 30602-7223
  - (706) 546-4238
- APGEN has been partially supported by the Georgia Research Alliance and has proprietary and patented technologies from UGA



## Why use phytoremediation?

- **≈** Plants are photosynthetic and have 80% of the energy in most ecosystems (the dominant organisms)
- **≈** Plants have extensive root systems for mining nutrients or pollutants (grow 100 million miles/acre/year).



APGEN, Inc 12/8/00 4

## Why use phytoremediation?

- ~ Remarkably effective.
- **≈**Cost effective 1/10<sup>th</sup> the expense of physical methods (capping, excavation, and reburial, soil roasting).
- **≫**Permanent solution as compared to reburial or capping of hazardous waste.
- **⊗**Green solutions improve the environment and have strong public support.

APGEN, Inc 12/8/00 5

## How does phytoremediation work?

- **≈**Organic pollutants are degraded to harmless small molecules (win-win).
- Elemental pollutants are detoxified, sequestered, and may be hyperaccumulated for harvest.



# APGEN's Approach to the Phytoremediation of Toxic Organics

- **≈**Treatability studies find particular plant species that degrade organic contaminants
- **≈** Plants selected from many native species environmentally suited to the site
- ≈Field test the plants and clean up the site



APGEN, Inc 12/8/00 7

APGEN's treatability studies identify those plant species that degrade organic pollutants in sediment or water samples from a site.





12/8/00

# APGEN's analytical methods quantify the degradation of pollutants by plants.





## We have quantitative data on the phytodegradation of organics such as:

- **≈**Chlorinated solvents
- **≫**Benzopyrenes
- **≈**Poly-Aromatic Hydrocarbons (PAHs)
- **≫**Phthalates
- **≫**Benzene
- &DDD
- **≫DDT**



APGEN, Inc 12/8/00 10

## Plant Selection for Solvent Waste Disposal Site


**≈**68 rapidly growing plant species suitable to the site were initially analyzed


≈ 14 species significantly accelerated the rate of degradation for one or more target compounds (methylene chloride, chloroform, benzene)




APGEN, Inc 12/8/00 11

# Selected plant species shorten the half-life of methylene chloride Sinoth Hill Species A B C D E F G H I J K L M N Control Plant Species AFGEN, Inc 12/8/00 12









Compact willow accelerates the rate of degradation for all three target compounds


APGEN, Inc 12/8/00 16



Red bay accelerates the rate of degradation for methylene chloride and chloroform, but not benzene

APGEN, Inc 12/8/00 17


## Selected plants dramatically enhance degradation of benzopyrenes from coal gassification site




30 species this range

3 species this range









Toxic elemental pollutants (e.g., heavy metals, radionucleides) present special problems

- **Elemental pollutants are immutable** short of nuclear fission or fusion
- ➢Phytoremediation goals are detoxification, sequestration, extraction, and/or hyperaccumulation




APGEN, Inc 12/8/00 22

# APGEN's Approach to Toxic Elemental Pollutants

- Test assertive genes in model plant species for enhancement of detoxification and/or accumulation
- **∞**Move genes into native conservation plant species suited to a variety of sites





"Mercury species are subject to much faster atmospheric removal than elemental mercury"

Keating *et al.*, 1997 EPA's "Mercury Study Report to Congress"

Conclusion: Hg(0) will be globally diluted into an ancient pool of mercury

"Mercury species are subject to much faster atmospheric removal than elemental mercury"

Keating *et al.*, 1997 EPA's "Mercury Study Report to Congress"

Conclusion from report: Hg(0) will be globally diluted into an ancient pool of mercury




APGEN, Inc 12/8/00 26

# Methylmercury is a more serious problem than ionic mercury.

- **≫**Methylmercury is more toxic
- **∞**Methylmercury is very efficiently biomagnified up the food chain up to 10<sup>8</sup> fold.
- Methylmercury is produced at all aquatic mercury sites



## Methylmercury can be broken down to less toxic non-biomagnified products





APGEN, Inc 12/8/00 28

First consider the MerA catalyzed reduction of ionic to less toxic metallic mercury


Hg(II) MerA Hg(0) ↑

Ionic mercury Metallic mercury





APGEN, Inc 12/8/00 29


Plants expressing the bacterial merA gene are resistant to levels of mercuric ion that kill controls.



50 μM Hg(II)








# Rice plants expressing merA clean Hg(II) from aquatic sediments





Inc 12/8/00 34



# Methylmercury is a more serious problem than ionic mercury.

- **≫**Methylmercury is more toxic
- $\gg$  Methylmercury is very efficiently biomagnified up the food chain up to  $10^8$  fold.
- **∞**Methylmercury is produced at all aquatic mercury sites



Minamata Bay, Japan and several other mercury poisoning incidents illustrated the tragic human consequences of methylmercury biomagnification in aquatic ecosystems.



APGEN, Inc 12/8/00 37

## Methylmercury can be broken down to less toxic non-biomagnified products

 $10^6 - 10^8$ 

Not

Not

Biomagnified

biomagnified

biomagnified

$$CH_3Hg^+ \xrightarrow{MerB} Hg(II) \xrightarrow{MerA} Hg(0)$$




APGEN, Inc 12/8/00 38


## APGEN's plants can handle methylmercury (MeHg) in several ways


- **≈**Detoxify MeHg and accumulate Hg(II) below ground
- **Detoxify MeHg and eliminate mercury from** site as Hg(0)
- **≈**Current research on detoxifying MeHg and hyperaccumulating Hg(II) above ground



Transgenic *merB* expression in *Arabidopsis* degrade methylmercury and accumulate Hg(II)







## **Summary of APGEN's Technologies**

- **≫**Native plants degrade organics efficientlyshould be no issues of public acceptance
- ➢ Heavy metals require transgenic technologies focusing on worst heavy metal toxins now and radionucleides in the future



#### **Planned Future Directions**

- •Engineering above-ground hyperaccumulation
- •Detoxification of other toxic heavy metal ions (cadmium, arsenic, copper, lead, & chromium)
- •Phytomining of gold, silver, & platinum
- •Genomics to identify thousands of relevant plant genes for enhancing phytoremediation


APGEN, Inc 12/8/00 43

Engineering a hyperaccumulator:
Wild-type shoots accumulate 10x more Hg(II)
when grafted with merA roots

wt

merA
wild-type (wt) grafted merA

Phytomining: The expression of *merA* confers gold ion (Au(III)) resistance to plants



#### Special Concerns for Peconic River Site #1

- **∞**Only native plant species used for organics
- > Wetlands viewed as a productive resource
- **≈**APGENs phytoremediation looks at most active species not just any species
- **≫**Native trees, herbaceous plants, and sterile grasses for transgenics



APGEN, Inc 12/8/00 46

## Special Concerns for Peconic River Site #2

- **≈** Have technology for Hg and DDD in place
- ≈Preliminary research on Ag and Cu.
- **≈** Treatability study could identify PCB degrading plants in 4-6 months
- **≈**Hg research supported by DOE EMSP
- &We're linked with CRADA for Peconic Site
  - Collaboration with Mark Furhmann (BNL), a phytoremediation & metals expert

